

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/289530644>

THE PLANT DESTROYER GENUS PHYTOPHTHORA IN THE 21ST CENTURY

Article · January 2014

CITATIONS

16

READS

475

2 authors, including:



[Sanjoy Guha Roy](#)

West Bengal State University

38 PUBLICATIONS 771 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Developing diagnostic parameters for control of Phytophthora infestans by studying genetic diversity using phenotypic, molecular and next gen sequencing methods. [View project](#)



Evaluation of genotypic diversity of field isolates of Phytophthora melonis causing fruit and vine rot of pointed gourd in West Bengal for formulation of effective control measures. [View project](#)

Rev. Plant Pathol. Vol. 6, 2014
Indian Society of Mycology and Plant Pathology
Scientific Publishers (India), Jodhpur pp. 387-412

THE PLANT DESTROYER GENUS PHYTOPHTHORA IN THE 21ST CENTURY

S. Guha Roy¹ and N.J. Grünwald²

¹Department of Botany, West Bengal State University, Barasat,
 Kolkata - 700126, WB, India

²USDA Agricultural Research Service, Corvallis, OR, USA

KEYWORDS: Aquatic Phytophthoras, Crop/Plantation Phytophthoras,
 Forest Phytophthoras integrated morphological & phylogenetic
 key, multi-host species, putative new species, genome
 sequencing

ABSTRACT

The 21st century has already seen a major paradigm shift in our understanding of the biology, evolution, and genetics of the genus *Phytophthora* as well as the tools and approaches used to develop novel approaches for disease management. The combination of novel tools and approaches provided by the convergence of genome sequencing, computing power and novel genomic/biotechnological tools paints a promising picture of the future of *Phytophthora* disease management. At the same time *Phytophthora* pathogens continue to emerge at an accelerated rate due to increased global travel and trade. This article provides a brief review of the impact, biology and the current genome-enabled era that will characterize progress in the 21st century.

INTRODUCTION

The genus *Phytophthora* has a long history in modern science. The scientific discipline of plant pathology was born in the early 1860s when Anton de Bary recognized *Phytophthora infestans* as the pathogen causing potato late blight responsible for the Irish potato famine in the 1840s (4,5). It also brought about the first formulated use of a fungicide. In addition to this substantial social and historical impact, even considering merely a handful of *Phytophthora* spp. (e.g., *P. sojae*, *P. infestans*) documented to cause significant economic impacts, costs amount to anywhere between 2-7 billions of dollars per crop per year worldwide in combined crop losses and management costs not even considering the less quantifiable but equally large impacts to natural ecosystems severely affected by some species (e.g., *P. cinnamomi*, *P. ramorum*).

TAXONOMY AND BIOLOGY

The genus *Phytophthora* is presently placed in the kingdom Straminipila (7), under the phylum Heterokonta, sub-phylum Pernosporomycotina, class Pernosporomycetes (Oomycetes), subclass Pernosporomycetidae, order Pythiales and family Pythiaceae. The taxonomy of *Phytophthora* has undergone an evolution in the way it has been studied; from the era of six morphospecies groups (119, 95, 109) to the era of phylogenetic clades (23) and ITS fingerprinting based keys (37) and now to a new era of an integrated morphological and phylogenetic key (102).

Validation of new *Phytophthora* taxa in recent years has been supported with phylogenies generated by sequence analysis of the internal transcribed spacer region (ITS) of the ribosomal DNA gene repeat, the translation elongation factor 1- α , β -tubulin and mitochondrial encoded cytochrome oxidase (cox) I, II, and spacer regions (23, 74, 86, 45). The multilocus phylogeny constructed for 82 *Phytophthora* species with seven molecular markers (8) confirmed the presence of the 10 well-supported clades for the genus previously observed (23). Until 2010, 108 recognized *Phytophthora* taxa have been recognized (94), but many more have been described since then, perhaps an indication of the furious pace of discovery of *Phytophthora* species in recent times (Table 1).

Table 1: Newly discovered/re-described species of *Phytophthora* in the last decade

Year	Name of species	Remarks	Reference
2001	<i>P. tropicalis</i> Aragaki & Uchida	formerly referred to as <i>P. palmivora</i> MF4 [= <i>P. capsici</i>]	5
	<i>P. oryzo-bladis</i> Wang and Lu (ex H H Ho)		57
	<i>P. pistaceae</i> Mirab.	formerly referred to as <i>P. megasperma</i> s. l. [sensu lato]	90
	<i>P. ramorum</i> Werres, de Cock & Man In't veld		120
2002	<i>P. europaea</i> Hansen & Jung		71
	<i>P. psychrophila</i> Jung & Hansen		71
	<i>P. uliginosa</i> Jung & Hansen		71
	<i>P. brassicae</i> de Cock & Man in 't Veld	formerly referred to as <i>P. porri</i>	81
	<i>P. ipomoeae</i> Flier & Grünwald		35
2003	<i>P. inundata</i> Brasier, Sanchez- Hernandez & Kirk	formerly referred to as <i>P. sp.</i> O-group	15
	<i>P. pseudosyringae</i> Jung & Delatour		72
	<i>P. nemorosa</i> Hansen & Reeser		54
2004	<i>P. alni</i> with subsp. <i>alni</i> , <i>multiformis</i> , and <i>uniformis</i> Brasier & Kirk	formerly referred to as alder <i>Phytophthora</i> , a species hybrid [Brasier et al., 1999]	13
	<i>P. hedraiandra</i> de Cock & Man in't Veld		26

2005	<i>P. kernoviae</i> Brasier, Beales & Kirk	formerly referred to as <i>P. taxon C</i>	14
2006	<i>P. captiosa</i> Dick & Dobbie		27
	<i>P. fallax</i> Dick & Dobbie		27
	<i>P. foliorum</i> Donahoo & Lamour		29
	<i>P. polonica</i> Belbahri, Melgarejo & Leford		7
2007	<i>P. alticola</i> Maseko, Coutinho & M.J. Wingf.		88
	<i>P. austrocedrae</i> Gresl. & Hansen		42
	<i>P. frigida</i> Maseko, Coutinho & MJ Wingf		88
	<i>P. siskiyouensis</i> Reeser & EM Hansen		101
	<i>P. rubi</i> (Wilcox and Duncan) Man in 't Veld, comb. nov.	formerly referred to as <i>P. fragariae</i> var. <i>rubi</i>	81
2008	<i>P. bisheria</i> Abad ZG, Abad JA & Louws		2
	<i>P. asparagi</i>	formerly referred to as <i>P. megasperma</i> s. l., <i>P. taxon asparagi</i>	105
	<i>P. gallica</i> T Jung & J Nechwatal	formerly referred to as <i>P. taxon G</i>	68
	<i>P. irrigata</i> C Hong & M Gallegly	formerly referred to as <i>P. drechsleri</i> s. l. [Dre I subgroup]	58
	<i>P. parsiana</i> Mostowfizadeh, Cooke & Banihashemi		91

	<i>P. pinifolia</i> AJ. Durán, Gryzenh. & MJ Wingf	formerly referred to as <i>P. cryptogea</i> or <i>P. drechsleri</i>	31
	<i>P. quercetorum</i> Balci & Balci		6
2009	<i>P. menzei</i> G.T. Browne, M.E. Gallegly & C.X. Hong	formerly referred to as <i>P. citricola</i> s. l. [avocado subgroup]	59
	<i>P. multivora</i> PM Scott & T Jung	formerly referred to as <i>P. citricola</i> s. l. / ' <i>P. citricola</i> complex'	108
	<i>P. plurivora</i> T. Jung and T.I. Burgess	formerly referred to as <i>P. citricola</i> s. l. / ' <i>P. citricola</i> complex' or <i>P. inflata</i>	67
	<i>P. rosacearum</i> EM Hansen & Wilcox	formerly referred to as <i>P. megasperma</i> s. l. [AC subgroup]	55
	<i>P. sansomeana</i> EM Hansen & Reeser	formerly referred to as <i>P. megasperma</i> s. l. [DF subgroup]	55
	<i>Phytophthora X pelgrandis</i> W.F. Gerlach, Nirenberg & Graffenh		96
2010	<i>P. andina</i> Adler & Flier	formerly referred to as <i>P. infestans</i> s. l.,	97
	<i>P. chrysanthemi</i> Naher M, Watanabe, H., Chikuo, Y., & Kageyama		93
	<i>P. hydropathica</i> Hong, C. and Gallegly, M.	Previously termed taxa Dre II. Closest relatives, <i>P. parsiana</i> and <i>P. irrigata</i> (Dre I). High temperature tolerant and isolated from irrigation water	61
	<i>P. morindae</i> Abad ZG & S Nelson	In ITS clade 10 and most closely related to <i>P. kernoviae</i> but differ in having umbellate sporangiophores. Host: Indian mulberry, <i>Morinda citrifolia</i> L. var. <i>citrifolia</i>	94

2011	<i>P. glovera</i> Z.G. Abad & H.D. Shew	In ITS clade 2 and most closely related to <i>P. capsici</i> but differs in being homothallic and produces both paragynous and amphigynous antheridia. Host :Tobacco	3
	<i>P. gibbosa</i> T. Jung, M.J.C. Stukely & T.I. Burgess, sp. nov.	In ITS clade 6 and associated with dying vegetation in swampy or water logged areas	69
	<i>P. gregata</i> T. Jung, M.J.C. Stukely & T.I. Burgess, sp. nov.		69
	<i>P. litoralis</i> T. Jung, M.J.C. Stukely & T.I. Burgess, sp. nov.	In clade 6 and present in water ways	69
	<i>P. thermophila</i> T. Jung, M.J.C. Stukely & T.I. Burgess, sp. nov.	In clade 6 and present in water ways; previously been misidentified as <i>P. drechsleri</i> and is referred to as <i>P. sp. 3</i> (Burgess et al. 2009)	69
	<i>P. himalsilva</i> Vettraino, Brasier & Vannini, sp. nov.	In ITS Clade 2 and forms part of a species cluster defined as Clade 2c by Blair et al. (2008)	116
	<i>P. pini</i> Leonian, 1925 Emend. Gallegly, Hong, Richardson & Kong = <i>Phytophthora citricola</i> I Gallegly & Hong 2008	Formerly known as <i>Phytophthora citricola</i> Cil I type (Kong et al. 2003, Gallegly and Hong 2008)	60
	<i>P. obscura</i> Grünwald & Werres, sp. Nov.	Defines a novel subclade 8d with close relatives <i>P. Syringae</i> and <i>P. Austrocedrae</i>	43

Although there have been considerable advances in the taxonomy of these organisms, the identification to species level using molecular characters remains a major challenge for plant pathologists. Several species "complexes" can be observed in

phylogenetic trees showing the presence of potential cryptic species. Presently, powerful molecular techniques combined with morphological characterisation and a renewed interest in probing of the environments have led to discovery of new species, novel variants within species, hybrids and provided a better resolution of species “complexes” with differentiation of the species therein (46).

There are numerous putative new species, which are in progress for official description in different laboratories around the world. These include *Phytophthora* (*P.*) taxa, such as *P. taxon* Pgchlamydo, *P. taxon* walnut, *P. taxon* oaksoil, *P. taxon* forestsoil, *P. taxon* riversoil, *P. taxon* salixsoil, *P. taxon* raspberry, *P. taxon* Banksia, *P. taxon* *meadii*-like, *P. taxon* chicory, *P. taxon* Acer, *P. taxon* Agathis, *P. taxon* orphan. Most of these originate from natural ecosystems and are shown to belong to the ITS clade 6 (7, 10, 23). Others are *P. taxon* paludosa (69), *P. sp.* personii (8) *P. taxon* rosacearum-like, *P. taxon* kwongan, *P. taxon* hungarica and *P. taxon* sulawesiensis. Many of these belong to clades 7 or 11 and are so far found only in Australian natural ecosystems where they are associated with plant mortalities (19). The discoveries of natural hybrids have added another dimension (9, 13, 82, 84, 38). And this is not the end; the following provisionally named species are awaiting their validation: *P. cuyabensis*, *P. hungarica*, *P. lagoariana*, *P. niederhauserii*, *P. parvasperma*, *P. sulawesiensis* and *P. sylvatica* (= *P. taxon* forestsoil?). Some, like *P. niederhauserii*, *P. asparagi*, *Phytophthora gemini* although not yet described formally, have been proposed as new species (4, 105, 82).

With the advent of molecular techniques, the most rapid resolution and differentiation of species has been in the ITS clades 2, 3, 6 & 7 (23). This, in some cases, has necessitated division into sub-clades; viz. clade 6, which previously consisted of only three species: *P. gonapodyides*, *P. humicola* and *P. megasperma* (33) has now radiated to the extent that it can be subdivided into 3 sub clades with 11 described species, many *P. taxons* and undescribed species (69). With description of more species in ITS clade 2, it has now been suggested that Clade 2c (8) has evolved within Asia (116).

Considering that there are 200- 600 extant *Phytophthora* species (10), a large number of species therefore remain to be discovered. Once more species are discovered and described,

clearer and well resolved species boundaries will emerge, resulting in revised phylogeny reconstructions. For example, it has become clear that *P. citricola* formerly recognized as a single taxon in phylogenetic reconstructions (23, 8) is now split into a group of species including *P. multivora* and *P. plurivora* (67). As more sequence data become available and new species are described, a clearer picture of species boundaries will emerge.

Host ranges of *Phytophthora* species can vary from one extreme of being very diverse (*P. ramorum*, *P. cinnamomi*, *P. nicotianae*) to that of a single host (*P. sojae*). Host ranges of new invasive forest or other *Phytophthora* species are being determined but the range continues to expand with passage of time (48, 107). Molecular validation and renaming of species are now often increasingly changing host boundaries for each species (e.g. *P. palmivora* MF4 [= *P. capsici*] - *P. tropicalis*). A comparative proteomics study between *P. ramorum* (broad host range) and *P. sojae* (single host range) identified some candidate proteins that might be factors involved in determination of host range (105), but more such comparisons are needed with other broad host range species whose genomes have been sequenced like *P. capsici* and *P. nicotianae* for validation and refinement. As host-pathogen (*A. thaliana*-*P. nicotianae*) interaction models for broad host range are also available (118), the focus perhaps needs to shift to multi-host species and finding the determinants for host specificity which would provide insights for control of these species which cause the most economic impact worldwide but have been difficult to control. Also, as proposed (89), species diverge by geographic isolation (allopatric) or, in the case of pathogens, by a switch in host (sympatric e.g. *P. infestans* - *Phytophthora mirabilis* - *Phytophthora ipomoeae*) and hence it is highly unlikely that two closely related species are able to diverge if they share a common host in the same geographical area. The study of host ranges will therefore become increasingly important in the future to answer questions of potential hosts of invasive pathogens and their speciation and emergence.

IMPACT

Phytophthora pathogens have a large impact on native ecosystems, forests and agricultural crops. The past decade has seen the discovery of a large number of *Phytophthoras* especially

from aquatic and forest ecosystems and a wealth of information is now available on some of the Phytophthoras attacking crops and plantations. These advances and their possibilities will only be discussed here as exhaustive reviews are already available on the subject (75, 51). Here, we selected a few major examples from three distinct habitats: Crop/plantations, forests, and aquatic environments.

Crop/Plantation Phytophthoras

For the most studied of all Phytophthoras, namely *Phytophthora infestans*, the population structure has changed dramatically in the 21st century due to migrations and establishment of sexual populations in select areas. In fact, at very few locations is the population structure similar to what it was in the 20th century. The populations in Mexico (which is thought to be the source of recent migrations) are sexual and this region represents the centre of origin of this pathogen (49, 35, 45). Populations in Asia, North America, South America and north of Mexico have changed and remain mostly clonal while population in parts of Europe have become sexual (36). Recent analysis of simple sequence repeats has revealed diversity within US clonal lineages (for e.g. US 22) (Fry, personal communication) and some new types have also been identified (92). Eucablight and now Euroblight are two websites of the concerted EU networks that rely on extensive multiplexed SSR analyses (78) for surveillance. These efforts have established that while 'blue 13' is a new virulent 'SSR genotype' found in most places in Europe, in France there has been a spatiotemporal change in population structures and a change in allele frequencies in the UK, suggesting a suspected gradual change in populations. A new set of multiplexed SSR primers (79) have revealed that some of the 'SSR genotypes' obtained by using the Eucablight primers could further be subdivided into more genotypes; the converse was also true (20). Targeting of different loci may bring about finer resolution of some of the SSR genotypes; however, if the resultant "groupings/clusters/SSR genotypes" is to be of practical field value towards control of the disease, then each of these SSR genotypes have to be correlated with phenotypic characters like mating type, fungicide sensitivity, growth temperature etc. Implementation of effective control strategies requires knowledge about the genetic

structure of population of plant pathogens (122), as control strategies must target a population instead of an individual if they are to be effective. Defining the genetic structure of a population is a logical first step in studies of fungal population genetics because the genetic structure of a population reflects its evolutionary history and its potential to evolve: aspects important for formulating disease management strategies. Genetic diversity analyses of populations at these locations have therefore become a priority.

P. capsici in many parts of the world are the most important factors limiting vegetable production (52, 75, 123). An excellent review on the population structures and host ranges in both the Americas and Korea on *P. capsici* is available (75). Interestingly the populations of *P. capsici* are very diverse in the US and clonal in neighbouring countries in contrast to the clonal nature of *P. infestans* found in the USA. *P. capsici* additionally affects plantation crops like bell pepper, betelvine and cocoa in countries like India and Vietnam. Populations in India have been found to have intraspecies variation depending on host and techniques used (21) while population diversity is lower in Vietnam (112). Cocoa isolates were however of a single clonal lineage in both countries.

P. nicotianae affects over 255 plant genera in 90 families (22) and is relatively neglected (by scientists and policy makers) considering that this species is one of the most common pathogens on ornamental plants, the cultivation and trade of which has been recognized as a principal pathway for the introduction and spread of invasive plant pathogens (cited in 80). Also it is a major pathogen of different vegetables in some tropical economies. The neglect is potentially influenced by having an extremely broad host range that makes estimation of economic impact difficult. The intraspecific variation of *P. nicotianae* was studied variously by RAPD [tobacco isolates, (124); vegetable and fruit isolates, (52)]; AFLP [floricultural hosts, (77) tobacco, (110)] and variable mitochondrial intergenic regions [various hosts, (80)]. The results were similar showing that intraspecific variability exists and in some cases were related to hosts. The future course of action for both *P. capsici* and *P. nicotianae* would be to find out the extent of the variability with special reference to host specificity.

P. sojae with a single host is considered to be a low risk pathogen because of limited outcrossing and low mutation and has been extensively reviewed (118). However, it is evident that pathotype structures, in clonal populations, are changing (30).

Forest *Phytophthoras*

There are many prominent forest *Phytophthora* species. Perhaps most notable is the sudden oak death epidemic caused by the pathogen *P. ramorum* (48, 104). This pathogen emerged simultaneously on rhododendrons in gardens, woodlands and nurseries in Germany and forests and nurseries in the USA. Sudden oak death is only found in the USA where coast live oak and tanoak forests are severely impacted. However, a recent significant epidemic on Japanese larch has been documented in the UK that is of global concern (12). In both Europe and North America the pathogen exhibits a very broad host range that makes control exceedingly difficult. *P. ramorum* is obviously an exotic pathogen that has been introduced into the US and Europe from elsewhere. Currently, three clonal lineages are being recognized that have been named after the continent (NA = North America; EU = Europe) on which they were first found (47). EU1 only affects Europe, while all three lineages are found in the US (40,39). It has also become evident that the EU1 clonal lineage was moved from Europe to the Pacific Northwest, either British Columbia, Canada or Washington, US, most likely via movement of ornamental plants such as *Rhododendron* spp. (39). The origin of this pathogen is not well established, but recent coalescent analysis determined that the three lineages are either reproductively isolated or geographically isolated even though both mating types exist and have diverged before the rise of modern agriculture (40). Several studies have monitored and described establishment, migration and clonal divergence within each of the continents on which *P. ramorum* is observed (40, 39, 115, 87, 99, 98). Despite the occasional presence of both mating types in nursery environments of the Western US, sexual reproduction has to date not been found (38).

Another prominent forest *Phytophthora* is *P. lateralis* affecting Port Orford Cedar in the Pacific North Western US (56). Like *P. ramorum* this pathogen is introduced to the US and has become a new pathogen on a new host. Unlike *P. ramorum*, *P.*

lateralis has only one known host. Recent work indicated that this pathogen originated in Asia (11). It is thought that *P. ramorum* might also have originated from Asia, but this remains to be established.

There exist numerous other forest *Phytophthora* species of which notable examples include *P. kernoviae* (14) and *P. alni* (13).

Aquatic *Phytophthoras*

Though *Phytophthora* species are commonly referred to as water moulds, little is known about their aquatic ecology and the species present in these habitats are poorly characterized relative to those present on crops. More than 25 species of *Phytophthora*, (including *P. ramorum*, the sudden oak death pathogen), have been isolated from irrigation reservoirs and natural waterways in riparian ecosystems and as well as from swampy or water-logged areas (17, 111, 121, 63,101, 58, 61, 69). A number of previously unknown taxa also have been documented in aquatic environments (15, 58, 68, 69). Systematic sampling efforts have been undertaken for early detection of these pathogens and because of the renewed interest in the assemblage of *Phytophthora* species already resident in streams in natural ecosystems: in western North America (101), south Eastern America (64, 65), South China (123) and Australia (69). However, such efforts are lacking in the Indian subcontinent.

Most of the species found in aquatic environments belong to clade 6 and show a strong association with both forest and riparian ecosystems and, with the exceptions of *P. taxon asparagi*, *P. gonapodyides*, *P. megasperma* and *P. rosacearum*, which have limited association with agriculture and horticulture. However all ITS Clade 6 members are not aquatic, such as *P. pinifolia*, *P. inundata*, *P. taxon PgChlamydo* and *P. gonapodyides* which can be opportunistic and sometimes aggressive tree pathogens (18, 31, 68, 70). The function of most of these taxa within the ecosystems is unclear. It has been hypothesized (15, 17) that a saprotrophic lifestyle and sexual degeneration for taxa in this clade may be connected to their aquatic habitat. Furthermore, due to the almost complete absence of heterothallism within Clade 6 and a lack of records of Clade 6 taxa from the tropics, it has been hypothesised (69) that a tropical origin seems unlikely. Whatever their origin

might be, the fact that recycling of water for irrigation is becoming increasingly important, baiting or filtering irrigation water for pathogenic *Phytophthora* species will increasingly become a useful monitoring tool in agricultural systems, including nurseries (63).

GENOME-ENABLED BIOLOGY

Three *Phytophthora* genomes have been sequenced to date and these include *P. sojae*, *P. ramorum* and *P. infestans* (114, 53). Several other genomes are considered, being or have been sequenced including *P. parasitica*, *P. cinnamomi* and *P. capsici* among others. Comparative genomics of these genomes has revealed a remarkable diversity of effectors that include avirulence genes now referred to as RxLR genes for a conserved amino acid motif characterizing this unique class of small secreted proteins.

Availability of whole genome sequences provides novel tools and approaches for research and translational applications of research. For example, discovery of mating type gene, metalaxyl resistance genes, and other functional and or adaptive traits is within grasp. With a novel class of functional genetic markers plant pathologists will be able to characterize *Phytophthora* populations more accurately and more rapidly, while breeders will be able to pyramid resistance genes more appropriately while using the discovered effect or repertoire for effect or-mediated R-gene discovery (43, 117).

CONCLUSIONS

Phytophthora biology currently is an area of work for plant pathologists undergoing rapid discovery of new species and brimming with the hope of novel, genome-enabled translational discoveries. The *Phytophthora* community is hoping to sequence the genome of each known *Phytophthora* species in a concentrated effort, which will provide a useful, novel platform for development of targeted disease management strategies and result in novel insights about the evolution and patterns of emergence of this important plant pathogen genus.

REFERENCES

1. Abad ZG and Coffey M 2008 Development of a Morphological/Phylogenetic Lucid Key for the identification of Oomycetes *Phytophthora* 3rd International *Phytophthora* and *Pythium* Workshop in Integration of Traditional and Modern Approaches for Investigating the Taxonomy and Evolution of *Phytophthora*, *Pythium* and Related Genera, (http://www.phytophthoradborg/pdf/3rd_Inter_workshoppdf)
2. Abad ZG, Abad JA, Coffey MD, Oudemans PV, Man in 't Veld WA, de Gruyter H, Cunnington J, and Louws FJ 2008 *Phytophthora bisheria* sp nov, a new species identified in isolates from the Rosaceous raspberry, rose and strawberry in three continents, *Mycologia* **100** 99–110
3. Abad ZG, Ivors KL, Gallup CA, Abad JA, and Shew DH 2011 Morphological and molecular characterization of *Phytophthora glovera* sp nov from tobacco in Brazil *Mycologia* **103** 341–350
4. Abad ZG, and Abad JA 2003 Advances in the integration of morphological and molecular characterization in the genus *Phytophthora* The case of *P. niederhauseria* sp Nov, *Phytopathology* **93** S1
5. Aragaki M, and Uchida JY 2001 Morphological distinctions between *P. capsici* and *P. tropicalis* sp Nov, *Mycologia* **93** 137–145
6. Balci Y, Balci S, Blair JE, Park S, Kang S, and McDonald WL 2008 *Phytophthora quercetorum* sp nov, a novel species isolated from eastern and north-central US oak forest soils, *Mycological Research* **11** 2906–916
7. Belbahri L, Moralejo E, Calmin G, Oszako T, Garcí'a JA, Descals E, and Lefort F 2006 *Phytophthora polonica*, a new species isolated from declining *Alnus glutinosa* stands in Poland, *FEMS Microbiology Letters* **26** 1165–174
8. Blair JA, Coffey MD, Park S, Geiser DM, and Kang S 2008 A multilocus phylogeny for *Phytophthora* utilizing markers derived from complete genome sequences, *Fungal Genetics and Biology* **45** 266–277
9. Bonants PJM, Hagenaar-de Weerd M, Man in 't Veld WA, and Baayen RP 2000 Molecular characterization of natural hybrids of *Phytophthora nicotianae* and *P. cactorum*, *Phytopathology* **90** 867–874
10. Brasier C 2009 *Phytophthora* biodiversity how many

- Phytophthora* species are there? in EM Goheen, SJ Frankel (tech coords) *Phytophthoras* in Forests and Natural Ecosystems Proc, 4th Meeting of IUFRO Working Party 070209, Monterey, CA Gen Tech Rep PSW-GTR-221, Albany, CA, 101–115 pages
11. Brasier CM, Vettraino AM, Chang TT, and Vannini A 2010 *Phytophthora lateralis* discovered in an old growth Chamaecyparis forest in Taiwan, *Plant Pathology* **59** 595-603
 12. Brasier C and Webber J 2010 Plant pathology Sudden larch death, *Nature* **466** 824-5
 13. Brasier CM, Kirk SA, Delcan J, Cooke DEL, Jung T, and Man in 't Veld WA 2004 *Phytophthora alni* sp nov and its variants designation of emerging heteroploid hybrid pathogens spreading on Alnus trees, *Mycological Research* **108** 1172–1184
 14. Brasier CM, Beales PA, Kirk SA, Denman S and Rose J 2005 *Phytophthora kernoviae* sp nov, an invasive pathogen causing bleeding stem lesions on forest trees and foliar necrosis of ornamentals in the UK, *Mycological Research* **109** 853–859
 15. Brasier CM, Sanchez-Hernandez E, and Kirk SA 2003a *Phytophthora inundata* sp nov, a part heterothallic pathogen of trees and shrubs in wet or flooded soils, *Mycological Research* **107** 477–484
 16. Brasier CM, Cooke DEL, and Duncan JM 1999 Origin of a new *Phytophthora* pathogen through interspecific hybridization, *PNAS, USA*, **96** 5878–5883
 17. Brasier CM, Cooke DEL, Duncan JM, and Hansen EM 2003b Multiple new phenotypic taxa from trees and riparian ecosystems in *Phytophthora gonapodyides*–*P. megasperma* ITS clade 6, which tend to be high-temperature tolerant and either inbreeding or sterile, *Mycological Research* **107** 277–290
 18. Brown AV and Brasier CM 2007 Colonization of tree xylem by *Phytophthora ramorum*, *P. kernoviae* and other *Phytophthora* species, *Phytopathology* **56** 227–241
 19. Burgess TI, Webster JL, Ciampini JA, White DW, Hardy GESTJ, and Stukely MJC 2009 Re-evaluation of *Phytophthora* species isolated during 30 years of vegetation health surveys in Western Australia using molecular techniques, *Plant Disease* **93** 215–223
 20. Chmielarz M, Sobkowiak S, Lebecka R, and Śliwka J 2010 Chosen characteristics

- of Polish *Phytophthora infestans* isolates PPO-Special Report no 14 39 – 44 Twelfth EuroBlight workshop Arras (France),
21. Chowdappa P, Brayford D, Smith J, and Flood J 2003 Molecular discrimination of *Phytophthora* isolates on cocoa and their relationship with coconut, black pepper and bell pepper isolates based on rDNA repeat and AFLP fingerprints, *Current Science* **84** 1235-1238
 22. Cline ET, Farr DF, and Rossman AY 2008 Synopsis of *Phytophthora* with accurate scientific names, host range, and geographic distribution Plant Health Progress doi101094/PHP-2008-0318-01-RS
 23. Cooke DEL, Drenth A, Duncan JM, Wagels G, and Brasier CM 2000 A molecular phylogeny of *Phytophthora* and related Oomycetes, *Fungal Genetics and Biology* **30** 17–32
 24. De Bary A 1873 The potato disease, *The Journal of the Quekett Microscopical Club* **31** 39-145
 25. De Bary A 1876 Researches into the nature of the potato fungus, *Phytophthora infestans*, *Journal of the Royal Agricultural Society of England Series* **2** 12149-154
 26. de Cock AWAM and Lévesque CA 2004 New species of *Pythium* and *Phytophthora*, *Studies in Mycology* **50** 481-487
 27. Dick MW, Dobbie K, Cooke DEL, and Brasier CM 2006 *Phytophthora captiosa* sp nov and *P. fallax* sp nov causing crown dieback of Eucalyptus in New Zealand, *Mycological Research* **110** 393 404
 28. Dick MW 2001 Straminipilous Fungi Kluwer Academic Publishers, The Netherlands
 29. Donahoo R, Blomquist CL, Thomas SL, Moulton JK, Cooke DEL, and Lamour KH 2006 *Phytophthora foliorum* sp nov, a new species causing leaf blight of azalea, *Mycological Research* **110** 1309–1322
 30. Dorrance A and Grünwald NJ 2009 *Phytophthora sojae* Diversity Among and Within Populations in Oomycete genetics and genomics diversity, interactions and genomic tools (eds) L Kurt and Sophien Kamoun John Wiley & Sons, 197-212 pages
 31. Durán A, Gryzenhout M, Slippers B, Ahumada R, Rotella A, Flores F, Wingfield BD, and Wingfield MJ 2008 *Phytophthora pinifolia* sp nov associated with a serious needle disease of *Pinus radiata* in Chile, *Plant Pathology* **57** 715–727

32. Durán A, Gryzenhout M, Drenth A, Slippers B, Ahumada R, Wingfield BD, and Wingfield MJ 2010 AFLP analysis reveals a clonal population of *Phytophthora pinifolia* in Chile, *Fungal Biology* **114** 746–752
33. Erwin DC and Ribeiro OK 1996 *Phytophthora* Diseases, Worldwide APS Press
34. Flier WG, Grünwald NJ, Kroon LPNM, Sturbaum AK, van den Bosch GBM, Garay-Serrano E, Fry WE, and Turkensteen LJ 2003 The population structure of *Phytophthora infestans* from the Toluca Valley in central Mexico suggests genetic differentiation between populations from cultivated potato and wild *Solanum* species, *Phytopathology* **93** 382-390
35. Flier WG, Grünwald NJ, Kroon LPNM, van den Bosch BM, Garay-Serrano E, Lozoya-Saldana H, Bonants PJM, and Turkensteen LJ 2002 *Phytophthora ipomoeae* sp nov, a new homothallic species causing leaf blight on *Ipomoea longipedunculata* in the Toluca Valley of central Mexico, *Mycological Research* **106** 848–856
36. Fry WE, Grünwald NJ, Cooke DE, McLeod A, Forbes GA, and Cao K 2009 Population Genetics and Population Diversity of *Phytophthora infestans* in Oomycete genetics and genomics diversity, interactions and genomic tools (eds) L Kurt and Sophien Kamoun John Wiley & Sons, 139-164 pages
37. Gallegly ME and Hong CX 2008 *Phytophthora* Identifying species by morphology and DNA fingerprints St Paul, Minnesota APS Press 158 page
38. Goss EM, Cardenas MC, Myers K, Forbes GA, Fry WE, Restrepo S, and Grünwald NJ 2011 The plant pathogen *Phytophthora andina* emerged via hybridization of an unknown *Phytophthora* species and the Irish potato famine pathogen, *P infestans*, *PLoS One* (in press)
39. Goss EM, Larsen M, Vercauteren A, Werres S, Heungens K, and Grünwald NJ 2011 *Phytophthora ramorum* in Canada Evidence for migration within North America and from Europe, *Phytopathology* **101** 166-171
40. Goss EM, Carbone I, and Grünwald NJ 2009 Ancient isolation and independent evolution of the three clonal lineages of the exotic sudden oak death pathogen *Phytophthora ramorum*, *Molecular Ecology* **18** 1161–1174

41. Goss EM, Larsen M, Chastagner GA, Givens DR, and Grünwald NJ 2009 Population genetic analysis infers migration pathways of *Phytophthora ramorum* in US nurseries, *PLoS Pathogens* **5**,e1000583
42. Greslebin AG, Hansen EM, and Sutton W 2007 *Phytophthora austrocedrae* sp nov, a new species associated with *Austrocedrus chilensis* mortality in Patagonia (Argentina), *Mycological Research* **111** 308–316
43. Grünwald NJ 2012 Genome sequences of *Phytophthora* enable translational plant disease management and accelerate research, *Canadian Journal of Plant Pathology*
44. Grünwald NJ, Martin FN, Larsen MM, Sullivan CM, Press CM, Coffey MD, Hansen EM, and Parke JL 2011 *Phytophthora*-IDorg A sequence-based *Phytophthora* identification tool, *Plant Disease* **95** 337-342
45. Grünwald NJ, Werres S, Goss EM, Taylor CR, and Fieland VJ 2011 *Phytophthora obscura* sp nov, a new species that is a member of the novel *Phytophthora* subclade 8d, *Plant Pathology*
46. Grünwald NJ and Goss EM 2011 Evolutionary and Population Genetics of Exotic and Re-emerging Pathogens Traditional and Novel Tools and Approaches, *Annual Review of Phytopathology* **49** 249-267
47. Grünwald NJ 2009 Standardizing the nomenclature for clonal lineages of the sudden oak death pathogen, *Phytophthora ramorum* *Phytopathology* **99** 792-795
48. Grünwald NJ, Goss EM, and Press CM 2008 *Phytophthora ramorum* a pathogen with a remarkably wide host-range causing sudden oak death on woody ornamentals, *Molecular Plant Pathology* **9** 729–740
49. Grünwald NJ and Flier WG 2005 The biology of *Phytophthora infestans* at its center of origin, *Annual Review of Phytopathology* **43** 171-190
50. Grünwald NJ, Flier WG, Sturbaum AK, Garay-Serrano E, van den Bosch TBM, Smart CD, Matuszak JM, Lozoya-Saldana H, Turkensteen LJ, and Fry WE 2001 Population structure of *Phytophthora infestans* in the Toluca Valley region of Central Mexico, *Phytopathology* **91** 882-890

51. Guha Roy S 2008 Progress in *Phytophthora* research identification, species diversity and population diversity, *Journal of Mycopathological Research* **46** 163 - 184
52. Guha Roy S, Bhattacharyya S, Mukherjee SK, and Khatua DC 2009 Molecular Identification of *Phytophthora* spp Affecting some Economically Important Crops in Eastern India through ITS-RFLP and Sequencing of the ITS Region, *Journal of Phytopathology* **157** 666-674
53. Haas BJ 2009 Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*, *Nature* **461** 393-398
54. Hansen EM, Reeser PW, Davidson JM, Garbelotto M, Ivors K, Douhan L, and Rizzo DM 2003 *Phytophthora nemorosa*, a new species causing cankers and leaf blight of forest trees in California and Oregon, USA, *Mycotaxon* **88** 129-138
55. Hansen EM, Wilcox WF, Reeser PW, and Sutton W 2009 *Phytophthora rosacearum* and *P sansomeana*, new species segregated from the *Phytophthora megasperma* "complex", *Mycologia* **101** 129-135
56. Hansen EM, Goheen DJ, Jules ES, and Ullian B 2000 Managing Port-Orford-Cedar and the introduced pathogen *Phytophthora lateralis*, *Plant Disease* **84** 4-10
57. Ho HH 2001 *Phytophthora oryzo-bladis*, a new name for *P fragariae* var *oryzo-bladis*, *Mycotaxon* **78** 17-21
58. Hong C, Gallegly ME, Richardson PA, Kong P, and Moorman GW 2008 *Phytophthora irrigata*, a new species isolated from irrigation reservoirs and rivers in eastern United States of America, *FEMS Microbiology Letters* **285** 203-211
59. Hong CX, Gallegly ME, Browne GT, Bhat RG, Richardson PA, and Kong P 2009 The avocado subgroup of *Phytophthora citricola* constitutes a distinct species, *Phytophthora menzei* sp Nov, *Mycologia* **101** 833-840
60. Hong CX, Gallegly ME, Richardson PA, and Kong P 2011 *Phytophthora pini* Leonian resurrected to distinct species status, *Mycologia* **103** 351-360
61. Hong CX, Gallegly ME, Richardson PA, Kong P, Moorman GW, Lea-Cox JD, and Ross DS 2010b *Phytophthora hydropathica*, a new pathogen identified from irrigation water, *Rhododendron catawbiense* and *Kalmia latifolia*, *Plant Pathology* **59** 913-921

62. Hong CX, Richardson PA, and Kong P 2008 Pathogenicity to ornamental plants of some existing species and new taxa of *Phytophthora* from irrigation water, *Plant Disease* **92** 1201-1207
63. Hong CX and Moorman GW 2005 Plant pathogens in irrigation water challenges and opportunities, *Critical Reviews in Plant Science* **24** 189-208
64. Hulvey J, Gobena D, Finley L, and Lamour K 2010 Co-occurrence and genotypic distribution of *Phytophthora* species recovered from watersheds and plant nurseries of eastern Tennessee, *Mycologia* **102** 1127-1133
65. Hwang J, Oak SW, and Jeffers SN 2009 Monitoring occurrence and distribution of *Phytophthora* species in forest streams in North Carolina using baiting and filtration methods In Goheen EM, Frankel SJ (tech coords), Proceedings of the fourth meeting of the International Union of Forest Research Organizations (IUFRO) Working Party S070209 Phytophthoras in forests and natural ecosystems Gen Tech Rep PSW-GTR-221 91- 95 US Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA, USA
66. Jiang RHJ, Tripathy S, Govers F, and Tyler BM 2008 RXLR effector reservoir in two *Phytophthora* species is dominated by a single rapidly evolving superfamily with more than 700 members *PNAS* **105(12)** 4874-4879
67. Jung T and Burgess TI 2009 Re-evaluation of *Phytophthora citricola* isolates from multiple woody hosts in Europe and North America reveals a new species, *Phytophthora plurivora* sp Nov, *Persoonia* **22** 95-110
68. Jung T and Nechwatal J 2008 *Phytophthora gallica* sp nov, a new species from rhizosphere soil of declining oak and reed stands in France and Germany, *Mycological Research* **112** 1195-1205
69. Jung T, Stukely MJC, Hardy GESJ, White D, Paap T, Dunstan WA, and Burgess TI 2011 Multiple new *Phytophthora* species from ITS Clade 6 associated with natural ecosystems in Australia evolutionary and ecological implications, *Persoonia* **26** 13-39
70. Jung T 2009 Beech decline in Central Europe driven by the interaction between *Phytophthora* infections and

- climatic extremes, *Forest Pathology* **39** 73-84
71. Jung T, Hansen EM, Winton L, Obwald W, and Delatour C 2002 Three new species of *Phytophthora* from European oak forests, *Mycological Research* **106** 397-411
72. Jung T, Nechwatal J, Cooke DEL, Hartmann GC, Blaschke M, Oswald WF, Duncan JM, and Delatour C 2003 *P pseudosyringae* sp nov, a new species causing root and collar rot of deciduous tree species in Europe, *Mycological Research* **107** 772-789
73. Kong P, Hong CX, Richardson PA, and Gallegly ME 2003 Single-strand-conformation polymorphism of ribosomal DNA for rapid species differentiation in genus *Phytophthora*, *Fungal Genetics and Biology* **39** 238-249
74. Kroon LP, Bakker FT, van den Bosch GB, Bonants PJ, and FlierWG 2004 Phylogenetic analysis of *Phytophthora* species based on mitochondrial and nuclear DNA sequences, *Fungal Genetics and Biology* **41** 766-782
75. Lamour K and Kamoun S 2009 Oomycete Genetics and Genomics Diversity, Interactions and Research Tools Wiley-Blackwell, 582 page
76. Lamour K 2009 *Phytophthora capsici* Sex, Selection, and the Wealth of Variation, in Oomycete genetics and genomics diversity, interactions and genomic tools (eds) L Kurt and Sophien Kamoun John Wiley & Sons, 165- 178 pages
77. Lamour KH, Daughtrey ML, BensonDM, Hwang J, and Hausbeck MK 2003 Etiology of *Phytophthora drechsleri* and *P nicotianae* (= *P parasitica*) diseases affecting floriculture crops, *Plant Disease* **87** 854-858
78. Lees AK, Wattier RAM, Shaw DS, Sullivan L, Williams NA, and Cooke DEL 2006 Novel microsatellite markers for the analysis of *Phytophthora infestans* populations, *Plant Pathology* **55** 311-319
79. Li Y, Govers F, Mendes O, Testa A, Jacobsen E, Huang SW, and van der lee TAJ 2010 A new set of highly informative SSR markers for *Phytophthora infestans* population analysis assembled into an efficient multiplex, *Molecular Ecology Resources* **10(6)** 1098-1105
80. Mammella MA, Cacciola SO, Martin F, and Schena L 2011 Genetic characterization of *Phytophthora nicotianae* by the analysis of polymorphic regions of the mitochondrial

- DNA, *Fungal Biology* **115** 432-442
81. Man in 't Veld WA, de Cock AWAM, Ilieva E, and Levesque CA 2002 Gene flow analysis of *Phytophthora porri* reveals a new species *Phytophthora brassicae* sp Nov, *European Journal of Plant Pathology* **108** 51-62
82. Man in 't Veld WA 2001 First report of natural hybrids of *Phytophthora nicotianae* and *P cactorum* on loquat in Taiwan, *Plant Disease* **85** 98-98
83. Man in't Veld WA 2007 Gen flow analysis demonstrates that *Phytophthora fragariae* var *rubi* constitutes a distinct species, *Phytophthora rubi* comb Nov, *Mycologia* **99** 222-226
84. Man in'tVeld WA, de Cock AWAM, and Summerbell RC 2007 Natural hybrids of resident and introduced *Phytophthora* species proliferating on multiple new hosts, *European Journal of Plant Pathology* **117** 25-33
85. Man in't Veld WA, Rosendahl KCHM, Brouwer H and De Cock AWAM 2011 *Phytophthora gemini* sp nov, a new species isolated from the halophilic plant *Zostera marina* in the Netherlands, *Fungal Biology* doi10.1016/j.funbio.2011.05.006
86. Martin FN and Tooley PW 2003 Phylogenetic relationships among *Phytophthora* species inferred from sequence analysis of mitochondrially encoded cytochrome oxidase I and II genes, *Mycologia* **95** 269-284
87. Mascheretti S, Croucher PJP, Vettraino A, Prospero S, and Garbelotto M 2008 Reconstruction of the Sudden Oak Death epidemic in California through microsatellite analysis of the pathogen *Phytophthora ramorum*, *Molecular Ecology* **172** 755-2768
88. Maseko B, Burgue's TI, Coutinho TA, and Wingfield MJ 2007 Two new *Phytophthora* species from South African Eucalyptus plantations, *Mycological Research* **111** 1321-1338
89. Mayr E 1942 Systematics and the Origin of Species, New York Columbia University Press
90. Mirabolfathy M, Cooke DEL, Duncan JM, Williams NA, Ershad D, and Alizadeh A 2001 *Phytophthora pistaciae* sp nov and *P melonis* the principal causes of pistachio gummosis in Iran, *Mycological Research* **105** 1166-1175
91. Mostowfizadeh-Ghalamfarsa R, Cooke DE, and Banihashemi Z 2008 *Phytophthora parsiana* sp nov, a new high temperature tolerant species,

- Mycological Research* **112** 783–794
92. Myers K, Small I, Jensen S, Zuluaga P, Guha Roy S, and Fry WE 2010 Characterization of *Phytophthora infestans* isolates from potato/tomato in 2010, *Phytopathology* **101** 261–262
93. Naher M, Motohash K, Watanabe H, Chikuo Y, Senda M, Suga H, Brasier C, and Kageyama K 2010 *Phytophthora chrysanthemi* sp nov, a new species causing root rot of chrysanthemum in Japan, *Mycological Progress*
94. Nelson S and Abad ZG 2010 *Phytophthora morindae*, a new species causing black flag on Noni (*Morinda citrifolia* L) in Hawaii, *Mycologia* **102** 122–134
95. Newhook FJ, Waterhouse FM, and Stamps DJ 1978 Tabular key to the species of *Phytophthora* de Bary Commonwealth, *Mycol Inst Mycol Pap No 3*, 20 page
96. Nirenberg HI, Gerlach WF, and Graffenhahn T 2009 *Phytophthora X p elgrandis*, a new natural hybrid pathogenic to *Pelargonium grandiflorum* hort, *Mycologia* **101** 220–231
97. Oliva RF, Kroon LPNM, Chaco'n G, Flier WG, Ristaino JB, and Forbes GA 2010 *Phytophthora andina* sp nov, a newly identified heterothallic pathogen of solanaceous hosts in the Andean highlands, *Plant Pathology* **59** 613–625
98. Prospero S, Grünwald NJ, Winton LM, and Hansen EM 2009 Migration patterns of the emerging plant pathogen *Phytophthora ramorum* on the west coast of the United States of America, *Phytopathology* **99** 739–749
99. Prospero S, Hansen EM, Grünwald NJ, and Winton LM 2007 Population dynamics of the sudden oak death pathogen *Phytophthora ramorum* in Oregon from 2001 to 2004, *Molecular Ecology* **16** 2958–2973
100. Reeser PW, Hansen EM, and Sutton W 2007 *Phytophthora siskiyouensis*, a new species from soil, water, myrtlewood (*Umbellularia californica*) and tanoak (*Lithocarpus densiflorus*) in southwestern Oregon, *Mycologia* **99** 639–643
101. Reeser PW, Sutton W, Hansen EM, Remigi P, and Adams GC 2011 *Phytophthora* species in forest streams in Oregon and Alaska, *Mycologia* **103** 22–35
102. Ristaino JB 2011 Key for Identification of Common *Phytophthora* Species, APS Press, St Paul, MN
103. Rizzo DM, Garbelotto M, Davidson JM, Slaughter GW, and Koike ST 2002 *Phytophthora ramorum* as

- the cause of extensive mortality of *Quercus* spp and *Lithocarpus densiflorus* in California, *Plant Disease* **86** 205–214
104. Rizzo DM, Garbelotto M, and Hansen EM 2005 *Phytophthora ramorum* Integrative research and management of an emerging pathogen in California and Oregon forests, *Annual Review of Phytopathology* **43** 309–335
105. Saude C, Hurtado-Gonzales O P, Lamour KH, and Hausbeck MK 2008 Occurrence of a *Phytophthora* sp pathogenic to asparagus (*Asparagus officinalis*) in Michigan, *Phytopathology* **98** 1075–1083
106. Savidor A, Donahoo RS, Hurtado-Gonzales O, Land ML, Shah MB, Lamour KH, and McDonald WH 2008 Cross-species Global Proteomics Reveals Conserved and Unique Processes in *Phytophthora sojae* and *Phytophthora ramorum*, *Molecular Cellular Proteomics* **78** 1501–1516
107. Schwingle BW and Blanchette RA 2008 Host range investigations of new, undescribed, and common *Phytophthora* spp isolated from ornamental nurseries in Minnesota, *Plant Disease* **92** 642–647
108. Scott PM, Burgess TI, Barber PA, Shearer BL, Stukely MJC, Hardy GESTJ, and Jung T 2009 *Phytophthora multivora* sp nov, a new species recovered from declining Eucalyptus, Banksia, Agonis and other plant species in Western Australia, *Persoonia* 221–13
109. Stamps DJ, Waterhouse GM, Newhook FJ, and Hall GS 1990 Revised tabular key to the species of *Phytophthora* Mycological Papers 162 CAB International Mycological Institute, Kew, Surrey
110. Sullivan, M J, Parks EJ, Cubeta MA, Gallup CA, Melton T A, Moyer JW, and Shew HD 2010 An assessment of the genetic diversity in a field population of *Phytophthora nicotianae* with a changing race structure, *Plant Disease* **94** 455–460
111. Tjosvold SA, Chambers DL, Koike ST, and Mori SR 2008 Disease on nursery stock as affected by environmental factors and seasonal inoculum levels of *Phytophthora ramorum* in stream water used for irrigation, *Plant Disease* **92** 566–573
112. Truong NV, Liew ECY, and Burgess LW 2010 Characterisation of *Phytophthora capsici* isolates from black pepper in

- Vietnam, *Fungal Biology* **114** (2-3) 160-170
113. Tyler BM 2007 *Phytophthora sojae* root rot pathogen of soybean and model oomycete, *Molecular Plant Pathology* **8** 1-8
114. Tyler BM 2006 *Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis, *Science* **313** 1261-1266
115. Vercauteren A, De Dobbelaere I, Grünwald NJ, Bonants P, Van Bockstaele E, Maes M, and Heungens K 2010 Clonal expansion of the Belgian *Phytophthora ramorum* populations based on new microsatellite markers, *Molecular Ecology* **19** 92-107
116. Vettriano AM, Brasier CM, Brown AV, and Vannini A 2011 *Phytophthora himalsilva* sp nov an unusually phenotypically variable species from a remote forest in Nepal, *Fungal Biology* **115** 275-287
117. Vleeshouwers V, Rietman H, Krenek P, Champouret N, Young C, Oh SK, Wang MQ, Bouwmeester K, Vosman B, Visser RGF, Jacobsen E, Govers F, Kamoun S, and Van der Vossen EAG 2008 Effector genomics accelerates discovery and functional profiling of potato disease resistance and *Phytophthora infestans* avirulence genes, *PLoS ONE* **3** e2875
118. Wang Y, Meng Y, Zhang M, Tong X, Wang Q, Sun Y, Quan J, Govers F, and Shan W 2011 Infection of *Arabidopsis thaliana* by *Phytophthora parasitica* and identification of variation in host specificity, *Molecular Plant Pathology* **12** 187-201
119. Waterhouse GM 1963 Key to the species of *Phytophthora* de Bary Mycol Pap, No 92, 1-22 CMI, Kew, UK
120. Werres S, Marwitz R, Man in 't Veld WA, de Cock AWAM, Bonants PJM, de Weerd M, Themann K, Ilieva E, and Baayen RP 2001 *Phytophthora ramorum* sp nov, a new pathogen on *Rhododendron* and *Viburnum*, *Mycological Research* **105** 1155-1165
121. Werres S, Wagner S, Brand T, Kaminski K, and Seipp D 2007 Survival of *Phytophthora ramorum* in recirculating irrigation water and subsequent infection of *Rhododendron* and *Viburnum*, *Plant Disease* **91** 1034-1044
122. Wolfe MS and Caten CE 1987 Populations of Plant Pathogens Their Dynamics and Genetics (eds), Blackwell Scientific Publications, Oxford UK
123. Zeng HC, Ho HH, and Zheng FC 2009 A survey of *Phytophthora* species on

- Hainan Island of South China, *Journal of Phytopathology* **157** 33-39
124. Zhang XG, Sun WX, Guo L, Yu JF, and Chang CJ 2003 Genetic and pathogenic variation among tobacco

black shank strains of *Phytophthora parasitica* var *nicotianae* from the main tobacco growing in China, *Journal of Phytopathology* **151** 259-266.